

Arabidopsis Blue Light Receptor Phototropin 1 Undergoes Blue Light-Induced Activation in Membrane Microdomains

Yiqun Xue^{1,2,8}, Jingjing Xing^{1,2,8}, Yinglang Wan^{1,8}, Xueqin Lv¹, Lusheng Fan³, Yongdeng Zhang⁴, Kai Song^{1,2}, Li Wang¹, Xiaohua Wang¹, Xin Deng¹, František Baluška⁵, John M. Christie⁶ and Jinxing Lin^{1,7,*}

¹Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China

²University of Chinese Academy of Sciences, Beijing 100049, China

³Department of Botany and Plant Sciences, University of California, Riverside, CA 92521, USA

⁴Department of Cell Biology, Yale University School of Medicine, New Haven, CT 06520, USA

⁵Institute of Cellular and Molecular Botany, University of Bonn, 53115 Bonn, Germany

⁶Institute of Molecular, Cell and Systems Biology, University of Glasgow, Bower Building, Glasgow G12 8QQ, UK

⁷College of Biological Sciences and Biotechnology, Beijing Forestry University, Beijing 100083, China

⁸These authors contributed equally to this article.

*Correspondence: Jinxing Lin (linjx@ibcas.ac.cn)

<https://doi.org/10.1016/j.molp.2018.04.003>

ABSTRACT

Phototropin (phot)-mediated signaling initiated by blue light (BL) plays a critical role in optimizing photosynthetic light capture at the plasma membrane (PM) in plants. However, the mechanisms underlying the regulation of phot activity at the PM in response to BL remain largely unclear. In this study, by single-particle tracking and stepwise photobleaching analysis of phot1-GFP proteins we demonstrated that in the dark phot1 proteins remain in an inactive state and mostly exist as monomers. Dimerization and the diffusion rate of phot1-GFP increased in a dose-dependent manner in response to BL. In contrast, BL did not affect the lateral diffusion of kinase-inactive phot1^{D806N}-GFP but did enhance its dimerization, suggesting that phot1 dimerization is independent of phosphorylation. Förster resonance energy transfer–fluorescence lifetime imaging microscopy analysis revealed that the interaction between phot1-GFP and a marker of sterol-rich lipid environments, AtRem1.3-mCherry, was enhanced with increased time of BL treatment. However, this BL-dependent interaction was not obvious in plants co-expressing phot1^{D806N}-GFP and AtRem1.3-mCherry, indicating that BL facilitates the translocation of functional phot1-GFP into AtRem1.3-labeled microdomains to activate phot-mediated signaling. Conversely, sterol depletion attenuated phot1-GFP dynamics, dimerization, and phosphorylation. Taken together, these results indicate that membrane microdomains act as organizing platforms essential for the proper function of activated phot1 at the PM.

Key words: phot1, VA-TIRFM, spatiotemporal dynamics, blue light signaling, membrane microdomains

Xue Y., Xing J., Wan Y., Lv X., Fan L., Zhang Y., Song K., Wang L., Wang X., Deng X., Baluška F., Christie J.M., and Lin J. (2018). *Arabidopsis* Blue Light Receptor Phototropin 1 Undergoes Blue Light-Induced Activation in Membrane Microdomains. *Mol. Plant*. **11**, 846–859.

INTRODUCTION

Phototropins (phot) play key roles in phototropism, chloroplast movement, stomatal opening, leaf expansion, and solar tracking in response to blue light (BL) (Christie, 2007). *Arabidopsis thaliana* has two phot, phot1 and phot2, with overlapping but not completely redundant physiological functions (Liscum and Briggs, 1995; Sakai et al., 2001; Christie, 2007; Mo et al., 2015).

Physiological processes mediated by phot1 and phot2 are therefore complex.

Studies aimed at understanding phot functions have so far focused on their individual signaling mechanisms, as well as their

Published by the Molecular Plant Shanghai Editorial Office in association with Cell Press, an imprint of Elsevier Inc., on behalf of CSPB and IPPE, SIBS, CAS.